

UNCLASSIFIED

AD NUMBER

AD120083

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors;
Administrative/Operational Use; 20 JAN 1956.
Other requests shall be referred to Office of Naval Research, Washington, DC 20360.

AUTHORITY

onr ltr 9 nov 1977

THIS PAGE IS UNCLASSIFIED

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

UNCLASSIFIED

AD 200083

Armed Services Technical Information Agency

Reproduced by

DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address: Armed Services Technical Information Agency, Document Service Center, Knott Building, Dayton 2, Ohio.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

77 No/20083

FILE COPY

Difference Methods on a Digital Computer
for Laplacian Boundary Value
and Eigenvalue Problems

GEORGE E. FORSYTHE

FC

Reprinted from

COMMUNICATIONS ON PURE AND APPLIED MATHEMATICS
Volume IX, No. 3

August, 1956

Copyright, 1955, Interscience Publishers, Inc. All rights reserved.

Difference Methods on a Digital Computer for Laplacian Boundary Value and Eigenvalue Problems*

GEORGE E. FORSYTHE
University of California, Los Angeles

1. Finite Difference Approximations

We begin with

DEFINITION 1. A simple closed curve $\Gamma: x(s) + iy(s)$ ($0 \leq s \leq s_t$) is said to be piecewise analytic if $x(s)$ and $y(s)$ are real analytic functions of the arc length s of Γ in each of a finite number t of closed intervals $0 = s_0 \leq s \leq s_1$, $s_1 \leq s \leq s_2$, \dots , $s_{t-1} \leq s \leq s_t$.

Let the open region R of the x, y -plane have a boundary C which is the union of a finite number of piecewise analytic simple closed curves. For real-valued functions $u = u(x, y)$ defined on $R \cup C$, denote by Δu the Laplacian $u_{xx} + u_{yy}$. In numerical computations R, C, u , and Δu are frequently replaced by corresponding quantities associated with a finite difference network. These may be introduced as follows (see [1] for a similar treatment):

For a given mesh constant $h > 0$ let a net consist of the lines $x = \mu h$, $y = \nu h$, $\mu, \nu = 0, \pm 1, \pm 2, \dots$. The points $(\mu h, \nu h)$ are called nodes. The M nodes which are within R form the net region R_h , assumed to be connectable by line segments of the net within R . A point $(\mu h, \nu h)$ of R_h is said to be a regular interior point if each of the four neighboring nodes $(\mu h \pm h, \nu h \pm h)$ is in $R \cup C$. All other points of R_h are called irregular interior points. Certain points P of R_h have some number p ($1 \leq p \leq 4$) of neighboring nodes P_1, \dots, P_p not in the open region R . For any such point P , let S_k denote the closed segment of the net joining P to P_k . For each $k = 1, \dots, p$, the point of $S_k \cap C$ closest to P will be called a boundary point of the net. The set of boundary points will be denoted by C_h . Some points of C_h may be nodes, but each irregular interior point of R_h has one or more neighboring points in C_h which are not nodes. (The definition of C_h given in [2], although intended to be the same, is not precise.)

*The research leading to this paper was sponsored jointly by the Office of Naval Research, project NR-044-144, and the Office of Ordnance Research, project TB2-0001(1210).

Corresponding to u , defined on $R \cup C$, is a real-valued net-function $u = u_h$ defined on $R_h \cup C_h$. Corresponding to Δu is $\Delta^{(h)}u$ defined as follows. For each point $P: (x, y)$ of R_h , let the neighboring points of $R_h \cup C_h$ be denoted by $(x+h_E, y)$, $(x, y+h_N)$, $(x-h_W, y)$, and $(x, y-h_S)$, where each of the h 's is positive ($E \sim$ "east"; $N \sim$ "north", etc.). Let $b_E = h_E/h$, $b_N = h_N/h$, etc. For regular interior points $b_E = b_N = \dots = 1$. Let $u(x, y) = u_P$, $u(x+h_E, y) = u_E$, $u(x, y+h_N) = u_N$, etc.

Define the *residual* r_P at P by

$$(1) \quad r_P = \frac{u_E}{b_E(b_E+b_W)B} + \frac{u_W}{b_W(b_E+b_W)B} + \frac{u_N}{b_N(b_N+b_S)B} + \frac{u_S}{b_S(b_N+b_S)B} - 4u_P,$$

where

$$B = \frac{1}{4b_N b_S} + \frac{1}{4b_E b_S}.$$

Then

$$(2) \quad \Delta^{(h)}u = r_P/h^2.$$

(The definition of $\Delta^{(h)}u$ in [2] differs by the factor $2B$.) In case P is a regular interior point, (1) reduces to the usual formula

$$(3) \quad r_P = u_E + u_W + u_N + u_S - 4u_P.$$

When there are irregular interior points, the operator $\Delta^{(h)}$ is not self-adjoint over R_h . For treating the eigenvalue problem it is often more convenient to deal with the self-adjoint part of $\Delta^{(h)}$, denoted here and in [2] by Δ_h .

2. Asymptotic Inequalities for All Eigenvalues

For the operator Δ , the eigenvalue problem

$$(4) \quad \begin{aligned} \Delta u + \lambda u &= 0 \quad \text{in } R, \\ u &= 0 \quad \text{on } C \end{aligned}$$

is known to have an infinite sequence of linearly independent eigenfunctions $u^{(k)}$ and corresponding eigenvalues $\lambda^{(k)}$, numbered so that

$$(5) \quad 0 < \lambda^{(1)} < \lambda^{(2)} \leq \lambda^{(3)} \leq \dots$$

Similarly, the eigenvalue problem

$$(6) \quad \begin{aligned} \Delta_h u + \lambda_h u &= 0 \quad \text{on } R_h, \\ u &= 0 \quad \text{on } C_h \end{aligned}$$

has a finite sequence of linearly independent eigenfunctions $u_h^{(k)}$ and eigenvalues $\lambda_h^{(k)}$, where

$$0 < \lambda_h^{(1)} < \lambda_h^{(2)} \leq \lambda_h^{(3)} \leq \dots \leq \lambda_h^{(M)}.$$

(Recall that M is the number of points of R_h .)

For rectangles whose sides are lines of the net, it is well known that

$$(7) \quad \lambda_h^{(k)} < \lambda^{(k)}, \quad k = 1, \dots, M;$$

see also below. Since for general regions R close lower bounds for the $\lambda^{(k)}$ are not easily found, the author has been investigating [3, 5a, 2, 4] whether (7) might also hold for other regions.

Suppose one deals with a polygonal region R which is a finite union of squares and half-squares of a net of mesh constant $2h$. Then R_{2h} has no irregular interior points, and neither does R_h , obtained by refining the net. At any point (x, y) of R_{2h} ,

$$(8) \quad \Delta_{2h} u = \Delta_h u + \frac{1}{4} h^2 L_h u,$$

where $L_h u = h^{-4} \{u_{xx\bar{x}\bar{x}} + u_{yy\bar{y}\bar{y}}\}$. Here

$$u_{xx\bar{x}\bar{x}} = u(x-2h, y) - 4u(x-h, y) + 6u(x, y) - 4u(x+h, y) + u(x+2h, y),$$

and $u_{yy\bar{y}\bar{y}}$ is defined analogously. Following the idea [3] of putting the fundamental eigenfunction $u_h = u_h^{(1)}$ of the problem for R_h into the minimization problem for R_{2h} , one finds that

$$(9) \quad \lambda_{2h}^{(1)} \leq \lambda_h^{(1)} - \frac{1}{4} h^2 \left(\sum_{(2h)} u_h L_h u_h \right) \left(\sum_{(2h)} u_h^2 \right)^{-1},$$

where $\sum_{(2h)}$ denotes a sum over the points of R_{2h} . Hence whenever

$$(10) \quad \sum_{(2h)} u_h L_h u_h > 0$$

it follows that

$$(11) \quad \lambda_{2h}^{(1)} < \lambda_h^{(1)}.$$

If (11) should hold also for h replaced by $h/2, h/4, h/8, \dots$, it would then follow that $\lambda_h^{(1)} < \lambda^{(1)}$, since the methods of [5] show that $\lambda_h^{(1)} \rightarrow \lambda^{(1)}$, as $h \rightarrow 0$.

Recent computation with SWAC (see Section 4 below) for the L -shaped region of [3] shows that (11) fails for $h = 1/16$. Hence (10) is not true for all polygonal regions of the type considered here. I conjecture, however, that (10) holds for any *convex* R which is the union of squares and half-squares of the net.

Lacking a proof of this conjecture, the author proved [3] an asymptotic form of (7) for $k = 1$, and in [5a] and [2] generalized it to include certain regions with curved boundaries (the case $k = 1$ of Theorem 2 below). Thus, under the hypotheses of Theorem 2 below, the desired inequality (7) has already been proved for the first eigenvalue, though only for sufficiently small h . Less delicate inequalities of type $\lambda_h^{(1)} < \lambda^{(1)}$, valid for all h , have since been found by Weinberger [6, 7] and Hersch [8] without such stringent hypotheses on R , by allowing R_h to correspond to a region a little larger than

R . Assuming $x(s)$, $y(s)$ defining Γ to be seven times continuously differentiable, Saul'ev [8a] has recently found inequalities of the type $|\lambda_h^{(k)} - \lambda^{(k)}| = O(h^{2-\epsilon})$, as $h \rightarrow 0$.

During the 1955 Berkeley Conference for Partial Differential Equations, at which this paper was presented, the author completed an extension of the results of [2] to all eigenvalues of Δ and Δ_h . This has been abstracted in [4], and will now be summarized. Let τ be the angle between the x -axis and the positive direction of the tangent to C . For any function v , let v_n denote the inner normal derivative of v on C . For any two functions v, w , define an inner product (v, w) by the formula

$$12(v, w) = \iint_R (v_{xx}w_{xx} + v_{yy}w_{yy}) dx dy + \int_C v_n w_n \sin^2 2\tau d\tau.$$

Let the infinite symmetric matrix D have elements $d_{ij} = (u^{(i)}, u^{(j)})$, $i, j = 1, 2, \dots$, where $u^{(k)}$ is an eigenfunction of (4) belonging to $\lambda^{(k)}$, normalized so that $\iint_R (u^{(k)})^2 dx dy = 1$. Corresponding to an m -fold multiple eigenvalue of Δ , we are to take m linearly independent functions $u^{(k)}$.

Define a sequence of real numbers $\gamma^{(k)}$, $k = 1, 2, \dots$, as follows. If $\lambda^{(k)}$ is a simple eigenvalue of (4), take $\gamma^{(k)} = d_{kk}$. If $\lambda^{(k)} = \lambda^{(k+1)} = \dots = \lambda^{(k+m-1)}$ is an m -fold multiple eigenvalue of (4), then define $\gamma^{(k)} \geq \gamma^{(k+1)} \geq \dots \geq \gamma^{(k+m-1)}$ to be the m (real) eigenvalues of (d_{ij}) , $i, j = k, \dots, k+m-1$, the corresponding m -rowed principal minor of D .

DEFINITION 2. A region R bounded by a finite number of piecewise analytic curves is said to have strictly convex corners if, at any point $x(s_j) + iy(s_j)$ where distinct analytic curves meet (see Definition 1), the interior angle is less than π (equality not permitted).

THEOREM 1. Let R , an open region with strictly convex corners, be bounded by C , the union of a finite number of piecewise analytic, simple closed curves. Define $h, \lambda^{(k)}, \lambda_h^{(k)}, \gamma^{(k)}$ as above. Then for $k = 1, 2, \dots$ one has $-\infty < \gamma^{(k)} < \infty$ and, as $h \rightarrow 0$,

$$(12) \quad \lambda_h^{(k)} \leq \lambda^{(k)} - \gamma^{(k)} h^2 + o(h^2).$$

A proof of Theorem 1 will appear in a later paper. If R is also convex, then C is a simple closed curve with $d\tau \geq 0$, and D is therefore a Gramian matrix. Hence all $\gamma^{(k)} > 0$, and we have the following corollary to Theorem 1:

THEOREM 2. Let R , a convex open region with strictly convex corners, be bounded by C , a piecewise analytic, simple closed curve. Then, with the above notation, $0 < \gamma^{(k)} < \infty$, and for each $k = 1, 2, \dots$ there exists $h_0(k) > 0$ such that $\lambda_h^{(k)} < \lambda^{(k)}$ if $h \leq h_0(k)$.

The case $h = 1$ of Theorems 1 and 2 is the result proved in [2]. (In Theorem 1 of [2] the assumption that R be simply connected was never used. Note: The proof of Lemma 1 in [2] is incorrect.)

As an asymptotic result Theorem 1 is the best possible, in the following sense. For a rectangle R of sides $\pi/p, \pi/q$, the relation (12) is actually an equality for all h , up to the term $o(h^2)$. For, corresponding to the eigenfunction $u^{(m,n)} = \sin mpx \sin nqy$ of both (4) and (6) (it is convenient here to use a double superscript m, n instead of the single superscript k), one has

$$\lambda^{(m,n)} = (mp)^2 + (nq)^2;$$

and, as $h \rightarrow 0$ over values of h for which R is a union of squares of the net,

$$\begin{aligned} \lambda_h^{(m,n)} &= \frac{\sin^2 (m\pi h/2) + \sin^2 (n\pi h/2)}{(h/2)^2} \\ (13) \quad &= \lambda^{(m,n)} - \frac{(mp)^4 + (nq)^4}{12} h^2 + o(h^2) \\ &= \lambda^{(m,n)} - (u^{(m,n)}, u^{(m,n)}) h^2 / 12 + o(h^2). \end{aligned}$$

Since $(u^{(m,n)}, u^{(\mu,\nu)}) = 0$ unless $|m-\mu| + |n-\nu| = 0$, one has $\gamma^{(m,n)} = (u^{(m,n)}, u^{(m,n)})/12$, even for multiple eigenvalues. Hence (13) shows that (12) is an equality for the rectangle R , up to the term $o(h^2)$.

3. Solving the Difference Equations on a Machine

Let us first consider solving the *Dirichlet difference problem*

$$(14a) \quad \Delta^{(h)} u = 0 \quad \text{on } R_h,$$

$$(14b) \quad u = \text{prescribed function on } C_h$$

on an automatic digital computer. Assume there are no irregular interior points of R_h , so that (14a) is equivalent to

$$(14c) \quad r_P = u_E + u_N + u_W + u_S - 4u_P = 0 \quad \text{on } R_h.$$

To solve the system of M equations (14c) in the unknown values of u at the M points of R_h , iterative methods are usually preferable to elimination methods, because of their simplicity. The *successive overrelaxation* process of [10] and [9] has proved very successful. In it the points of R_h are repeatedly scanned in a cyclic order. In its turn, at each point P of R_h the value u_P is immediately replaced by a new value u'_P ,

$$(15) \quad u'_P = u_P + \frac{1}{4}\beta r_P,$$

where β ($1 \leq \beta < 2$) is a parameter to be discussed below. Using the new value of u_P , one goes on to the next point of R_h , and so on. For $\beta = 1$ this

is the *Liebmann process* [11], in which $u'_P = \frac{1}{4}(u_E + u_N + u_W + u_S)$. Any reasonable initial values $u^{(0)}_\beta$ of the function u over R_h will suffice, and one may take all $u^{(0)}_\beta = 0$.

Let $u^{(k)}_\beta$ be the result of k sweeps through R_h with successive overrelaxation. Let $u^{(\infty)}$ be the exact solution of (14); define the *error* $e^{(k)}_\beta = u^{(k)}_\beta - u^{(\infty)}$. Since the solution process is linear, one knows that

$$e^{(k)}_\beta = H^k_\beta e^{(0)}_\beta,$$

where H_β is a certain linear transformation of R_h into itself. The convergence of $e^{(k)}_\beta$ to 0, as $k \rightarrow \infty$, depends on the maximum modulus of the eigenvalues $\lambda_i(H_\beta)$ of H_β . For the Dirichlet problem it is known [9] that $|\lambda_i(H_\beta)| < 1$ (all i , $0 < \beta < 2$), so that successive overrelaxation ($1 \leq \beta < 2$) always converges. The ultimate speed of convergence depends on $\max_i |\lambda_i(H_\beta)|$. Now, for any single step (15), to choose $\beta = 1$ is the best *tactic*, in the sense that the negative definite error function $\sum_{R_h} e \Delta^{(h)} e$ is brought as close to the desired value 0 as is possible in the one step, by choosing $\beta = 1$ in (15). However, hand computers have long observed [12] that certain choices $\beta > 1$ apparently represent a better *strategy* than $\beta = 1$, in the sense that $e^{(k)}_\beta \rightarrow 0$ faster asymptotically, as $k \rightarrow \infty$. (Related ideas of strategy and tactics for other linear processes will be found in [13].) In [10] and [9] the matter of a best strategy is settled by showing how to pick the unique β_0 for which $\max_i |\lambda_i(H_\beta)|$ is minimized. Indeed,

$$(16) \quad \beta_0 = \frac{2}{1 + \sqrt{1 - \sigma}},$$

where

$$(17) \quad \sigma = \max_i |\lambda_i(H_1)|.$$

The number σ is easily approximated by setting $\beta = 1$ and estimating the asymptotic value of $\|e^{(k+1)}_1\|/\|e^{(k)}_1\|$, as $k \rightarrow \infty$, where $\|\cdot\|$ denotes some convenient norm function. When $\beta = \beta_0$, all the $|\lambda_i(H_\beta)|$, $i = 1, \dots, M$, turn out to be equal, so that the manner of approach of $\|e^{(k)}_{\beta_0}\|$ to 0 is utterly unpredictable.

When R has a curvilinear boundary C , the theory of [9] still holds, and the successive overrelaxation method is just as attractive. One simply replaces the formula (3) for r_P at irregular interior points P of R_h by the formula (1).

For the eigenvalue problem (6) successive overrelaxation is readily adaptable to getting $\lambda^{(1)}_h$ and its associated eigenvector $u^{(1)}_h$. One uses a trial value λ for $\lambda^{(1)}_h$, and solves the system

$$(18) \quad \begin{aligned} (\Delta_h + \lambda I)u &= 0 & \text{on } R_h, \\ u &= 0 & \text{on } C_h \end{aligned}$$

by successive overrelaxation starting with some nonzero vector $u_\beta^{(0)}$. For the residual, assuming that R_h has only regular interior points (so that $\Delta_h \equiv \Delta^{(h)}$), one uses

$$(19) \quad r_P = u_E + u_N + u_W + u_S - 4u_P + \lambda u_P$$

in (15). If λ were exactly equal to $\lambda_h^{(1)}$, the iterates $u_\beta^{(k)}$ would stay roughly in the affine subspace S (of M -dimensional space) which is orthogonal to $u_h^{(1)}$ and passes through $u_\beta^{(0)}$. Apart from an exceptional set of $u_\beta^{(0)}$ of zero measure, one would always have $u_\beta^{(k)} \rightarrow \alpha u_h^{(1)}$, as $k \rightarrow \infty$, where α is some nonzero number depending on $u_\beta^{(0)}$. If λ is near $\lambda_h^{(1)}$ but not equal to it, one finds that $u_\beta^{(k)}$ moves asymptotically into the line $\alpha u_h^{(1)}$ while converging to 0 or ∞ . An improved value of λ is then obtainable from the Rayleigh quotient $\varrho_h(u_\beta^{(k)})$, where

$$(20) \quad \varrho_h(v) = - \left(\sum_{R_h} v \Delta_h v \right) \left(\sum_{R_h} v^2 \right)^{-1}.$$

One then uses the better λ in (19), and alternates a few steps of successive overrelaxation with the Rayleigh quotient calculation, until a satisfactory $\alpha u_h^{(1)}$ is obtained.

For getting $\lambda_h^{(k)}$ and $u_h^{(k)}$ ($k > 1$), the idea of solving (18) with λ near $\lambda_h^{(k)}$ is undoubtedly still a good one, but successive overrelaxation no longer converges, because $\Delta_h + \lambda_h^{(k)} I$ is not semidefinite for $k > 1$. A simple iteration which can solve indefinite systems must be substituted—perhaps that of Kaczmarz [14].

4. SWAC Codes

SWAC is an electronic digital computer built by the National Bureau of Standards, but now owned by the Department of Defense and loaned to the University of California, Los Angeles, for research in numerical analysis. It is a three or four address machine with a Williams tube memory of 256 words; each word consists of 36 binary digits and a sign. Add time is .000064 second; multiplication time is slightly less than six add times. An auxiliary magnetic drum memory holds 4096 words in 128 channels of 32 words each (increased to 8192 words in late 1955). One can transfer a block of up to 32 words between the drum and the Williams tube memory in .017 second. For substantial problems, computing time is measured in hours, while coding time is measured in weeks.

In connection with the author's graduate seminar on numerical methods for elliptic partial differential equations, held in the spring of 1955, a few general codes were prepared following the methods of Section 3 above.

In all these codes the net region R_h is represented by a subset of the

32 by 128 rectangular array of cells on SWAC's magnetic drum. Parameters indicate the smallest rectangle in which R_h is contained. In all the codes the data in successive drum channels are brought to the Williams tube memory for the relaxation process, and the improved values are sent back when they are no longer needed.

The author's basic Code 60 solves Dirichlet's problem for the operator $\Delta^{(h)} \equiv \Delta_h$ for a region with no irregular interior points. Here C_h consists solely of nodes of the net, and can thus also be represented by cells of the drum. Points of R_h have a 0 in the most significant binary digit. Points not in R_h are tagged with a 1 in the same digit, and can be an arbitrary subset of the 32 by 128 rectangular array. Code 60 assumes that the values of u have been placed in the drum cells representing C_h , in digits less significant than the first (which is assumed to be tagged with a 1, as stated above). It performs successive overrelaxation at all points of R_h , leaving all other values of u unchanged. The overrelaxation factor β defined in Section 3 can be modified at will by the SWAC operator. (It is fun for the operator to participate in the solution. Having the operator manage β , while SWAC manages all other numbers, seems a satisfactory division of labor!) The error measure used, denoted by $\|e_\beta^{(k)}\|$, is the first power norm of $r^{(k)}$; it is displayed at the end of each sweep through R_h . It is easily accumulated during the algorithm (15), since

$$\|e_\beta^{(k)}\| = \sum_{R_h} |r_P^{(k)}|.$$

The ratio $\|e_\beta^{(k)}\| / \|e_\beta^{(k-1)}\|$ is also displayed, and is useful, for example, in approximating the σ of (17).

When R_h is a rectangle of 30 by 68 interior points, a relaxation sweep through the 2040 points takes SWAC 8.5 seconds. When $\beta = 1$, to reduce $e_1^{(k)}$ by a factor of 10^{-6} would take about 2300 sweeps—over five hours on SWAC. Using the optimal value $\beta = \beta_0 = 1.858+$, only 90 sweeps would be needed—about 13 minutes on SWAC. Including the time necessary to estimate that 1.875 is a pretty good value for β , and running with this β , a reduction of 10^{-6} in $e_\beta^{(k)}$ is found in practice to take 20 or 30 minutes (see [15]).

Code 61 is a modification of Code 60 in which boundary conditions corresponding to $\partial u / \partial n = u_n = 0$ are permitted at any point of C_h . Suppose, for example, that it is desired to have $\partial u / \partial x = 0$ for $x = 0$, where R is a region with $x > 0$. Then values of $u(x, y)$ for the lines $x = -h$, $x = 0$, $x = h$, $x = 2h$, \dots are all stored on the drum for appropriate values of y . When the code comes to $u(-h, y)$ for any y , the value is simply replaced by $u(h, y)$. The values $u(0, y)$, $u(h, y)$, $u(2h, y)$, \dots are all relaxed according to (15). Again the relaxation routine is notified of the different types of interior and boundary points by suitable identification digits.

Mr. Kenneth Ralston of U.C.L.A. has prepared a SWAC code analogous to Code 60, dealing with curvilinear boundaries. Note in this case that it is no longer possible to map C_h onto the drum in a simple manner. For example, the point of C_h "north" of the point $(\mu h, \nu h)$ of R_h may not be the same as the point of C_h "west" of $(\mu h + h, \nu h + h)$. Identification digits distinguish: (a) points not in R_h , to be left alone, (b) regular interior points, to be treated by (15), with r_p as in (3), and (c) irregular interior points, to be treated by (15), with r_p as in (1).

To deal with irregular interior points Ralston stores the coefficients of u_E, u_W, u_N, u_S in formula (1) in four cells of certain special channels of the drum. Since one meets the irregular interior points of R_h in a fixed order, all these special coefficients can be stored once and for all in the proper sequence. (Such of the u_E, u_W, u_N, u_S as are fixed boundary values are actually multiplied into the coefficients stored in the special channels.) Ralston can also use the special channels to simulate a general mixed boundary condition $au_n + bu + c = 0$ on a straight boundary segment parallel to the x or y axis.

The author's Code 70 performs successive overrelaxation for the eigenvalue problem (6) over regions with no irregular interior points, following the method described in Section 3. The quantities β and $4 - \lambda$ are both parameters of the code. Auxiliary to Code 70 are three other codes: Code 80 doubles the values of u over R_h , while Code 82 halves these values of u . Code 81 forms the Rayleigh quotient (20).

With Codes 70–82 the author found the fundamental eigenvalue for the L -shaped region of [3] for various h , and found the counter-example mentioned after (11). Code 61 has been adapted for some soil engineering calculations by Mr. D. Isherwood and Mrs. Louise Straus of U.C.L.A. Code 60 has been modified by Miss Rita Powers at U.C.L.A. to study the influence of the order in which the points of R_h are relaxed on the rate of convergence of successive overrelaxation.

To make Ralston's code more available for problems, a code is badly needed to tag the drum cells with the identification digits, to precompute the coefficients (1), and to store the latter on the special channels. Ralston and the author have outlined such a code, assuming C to be a piecewise quadratic function of x, y . Another needed code is one to get $\lambda_h^{(k)}$, for $k > 1$.

Using the ORACLE computer at Oak Ridge National Laboratories, W. C. Sangren [unpublished] has also computed $\lambda_h^{(1)}$ for the L -shaped region of [3] and for other simple regions with rectilinear boundaries.

Bibliography

- [1] Kantorovič, L. V., and Krylov, V. I., *Približennye Metody Vysšego Analiza*, 4th Edit., Gostehizdat, Moscow-Leningrad, 1952.
- [2] Forsythe, G. E., *Asymptotic lower bounds for the fundamental frequency of convex membranes*, Pacific J. Math., Vol. 5, 1955, pp. 691-702.
- [3] Forsythe, G. E., *Asymptotic lower bounds for the frequencies of certain polygonal membranes*, Pacific J. Math., Vol. 4, 1954, pp. 467-480.
- [4] Forsythe, G. E., *Asymptotic lower bounds for all the eigenvalues of a convex membrane*, Bull. Amer. Math. Soc. Abstract, Vol. 61, 1955, p. 551.
- [5] Courant, R., Friedrichs, K. O., and Lewy, H., *Über die partiellen Differenzengleichungen der mathematischen Physik*, Math. Ann., Vol. 100, 1928, pp. 32-74.
- [5a] Forsythe, G. E., *Arbitrarily close lower bounds for the fundamental frequency of a convex membrane*, Bull. Amer. Math. Soc. Abstract, Vol. 59, 1953, p. 551.
- [6] Weinberger, H. F., *Lower bounds for minima by finite difference methods*, Bull. Amer. Math. Soc. Abstract, Vol. 61, 1955, p. 301.
- [7] Weinberger, H. F., *Upper and lower bounds for eigenvalues by finite difference methods*, Comm. Pure Appl. Math., Vol. 9, 1956 (this issue) pp. 000-000.
- [8] Hersch, J., *Équations différentielles et fonctions de cellules*, C. R. Acad. Sci. Paris, Vol. 240, 1955, pp. 1602-1604.
- [8a] Saul'ev, V. K., *K voprosu rešeniya zadaci o sobstvennykh značeniyaх metodom konečnykh raznostei*, Vyčisl. Mat. Vyčisl. Tehn., Vol. 2, 1955, pp. 116-144.
- [9] Young, D., *Iterative methods for solving partial difference equations of elliptic type*, Trans. Amer. Math. Soc., Vol. 76, 1954, pp. 92-111 (condensation of his Harvard dissertation 1950).
- [10] Frankel, S. P., *Convergence rates of iterative treatments of partial differential equations*, Math. Tables and Other Aids to Computation, Vol. 4, 1950, pp. 65-75.
- [11] Liebmann, H., *Die angenäherte Ermittlung harmonischer Funktionen und konformer Abbildung*, S.-B. Math.-Nat. Kl. Bayer. Akad. Wiss., Vol. 47, 1918, pp. 385-416.
- [12] Fox, L., *A short account of relaxation methods*, Quart. J. Mech. Appl. Math., Vol. 1, 1948, pp. 253-280.
- [13] Stiefel, E., *Relaxationsmethoden bester Strategie zur Lösung linearer Gleichungssysteme*, Comment. Math. Helv., Vol. 29, 1955, pp. 157-179.
- [14] Kaczmarz, S., *Angenäherte Auflösung von Systemen linearer Gleichungen*, Bull. Int. Acad. Polon. Sci. Cl. Sci. Math. Nat. Ser. A Sci. Math., 1937, pp. 355-357.
- [15] Forsythe, G. E., *Relaxation methods*, Section 125 of I. S. Sokolnikoff, *Mathematical Theory of Elasticity*, McGraw-Hill, New York, 1956.

Received January 20, 1956.